

Frequency-dependent polarizabilities of alkali atoms from ultraviolet through infrared spectral regions

M. S. Safronova and Bindiya Arora

Department of Physics and Astronomy, University of Delaware, Newark, Delaware 19716

Charles W. Clark

Physics Laboratory, National Institute of Standards and Technology, Technology Administration,

U.S. Department of Commerce, Gaithersburg, Maryland 20899-8410

(Dated: February 2, 2008)

We present results of first-principles calculations of the frequency-dependent polarizabilities of all alkali atoms for light in the wavelength range 300-1600 nm, with particular attention to wavelengths of common infrared lasers. We parameterize our results so that they can be extended accurately to arbitrary wavelengths above 800 nm. This work is motivated by recent experiments involving simultaneous optical trapping of two different alkali species. Our data can be used to predict the oscillation frequencies of optically-trapped atoms, and particularly the ratios of frequencies of different species held in the same trap. We identify wavelengths at which two different alkali atoms have the same oscillation frequency.

PACS numbers: 31.15.Md, 32.10.Dk, 32.70.Jz, 32.80.Rm

I. INTRODUCTION

The trapping of two different species of ultracold gases has been an active topic of research for about ten years (see [1] and references therein), and has led to the simultaneous production of quantum degenerate Bose-Einstein and Fermi-Dirac gases [2, 3, 4] and recently to the observation of heteronuclear Feshbach resonances involving Bose-Einstein and Fermi-Dirac species [5, 6].

Two recent experiments [5, 6] provide a glimpse of the possibilities of mapping out the phase diagram of a Bose-Einstein/Fermi-Dirac mixture, potentially a rich system that remains to be explored. Our present work provides accurate estimates of the trapping frequencies of disparate species confined in the same optical trap. We have been attentive to finding cases in which all species, Bose-Einstein and Fermi-Dirac atoms, and the molecules formed from them, have the same trapping frequencies. This case is of particular interest because it has been shown to be optimal for a recently proposed scheme of cooling mediated by Feshbach resonances [7]. In addition, we note that while the DC polarizabilities of alkali atoms have been subject to careful experimental investigation, there is relatively little accurate experimental data on frequency-dependent polarizabilities. We draw attention to the fact that current experiments offer the possibility of measurements of unprecedented accuracy regarding the ratio of frequency-dependent polarizabilities of different species. Such measurements could provide more decisive comparison of *ab initio* atomic theory with experiments, than has been made to date.

We begin with a summary of the calculation of frequency-dependent polarizabilities, the details of which have been presented in previous publications [8, 9]. The computational method we use here has delivered agreement with experiments on static polarizabilities at the 1% level. There are few direct measurements of

frequency-dependent polarizabilities - the experiments are much more demanding, and have larger uncertainties associated with modeling the intensity profile of a focused laser beam. However, the alkali metal atoms are a special case, at least in the infrared spectral region. There, the polarizability is dominated by the contribution from the longest-wavelength resonance transitions, and it is possible to perform consistency checks between static values and long-wavelength values of the frequency-dependent atomic polarizabilities. We present tabulated values of calculated frequency-dependent polarizabilities over a range of wavelengths from the ultraviolet through the infrared spectrum, with particular attention to those wavelengths (primarily in the infrared) which have been or might be employed in the present generation of optical traps. For the infrared region, we display simple formulae that allow accurate computation of frequency-dependent polarizabilities for wavelengths not explicitly reported upon here.

Finally, we enumerate selected wavelengths, at which two species of optically-trapped alkali atoms would have the same frequencies of oscillation in a common optical trap. Such wavelengths offer opportunities for mapping out phase diagrams of Bose-Einstein and Fermi-Dirac mixtures in a space of reduced dimensionality. Moreover, to the extent that molecular resonances consist of weakly bound states at large internuclear separation, the oscillation frequencies of trapped molecular species should be very close to the frequencies of their atomic components. This may present an ideal system for investigation.

II. CALCULATION OF POLARIZABILITIES

The frequency-dependent polarizability of an atom in its ground state may be separated into polarizability of the ionic core α_{core} and the valence contribution α_v^{ns} .

TABLE I: The values of the frequency-dependent polarizabilities of alkali-metal atoms in a_0^3 . The wavelengths (in air) are given in nm, the corresponding frequencies are given in a.u.

λ_{air}	ω	Li	Na	K	Rb	Cs	Fr
1554	0.02931	201.0(7)	189.4(2)	381.2(8)	424.0(7)	570.9(1.2)	407.9(2.0)
1550	0.02939	201.3(7)	189.6(2)	381.8(8)	424.7(7)	572.2(1.2)	408.5(2.0)
1540	0.02958	201.9(7)	190.0(2)	383.4(8)	426.6(7)	575.6(1.2)	410.1(2.0)
1340	0.03399	218.1(8)	201.0(2)	428.2(9)	479.5(7)	673.7(1.3)	455.0(2.1)
1332	0.03420	219.0(8)	201.6(2)	430.7(9)	482.5(7)	679.5(1.3)	457.5(2.1)
1313	0.03469	221.1(8)	203.0(2)	436.9(9)	489.9(7)	694.1(1.3)	463.7(2.1)
1240	0.03673	231.0(8)	209.3(2)	465.9(9)	524.6(7)	764.6(1.4)	492.8(2.2)
1152	0.03954	247.0(8)	219.4(2)	516.1(1.0)	585.7(8)	899.9(1.6)	543.6(2.3)
1090	0.04179	262.7(8)	228.8(2)	568.7(1.1)	650.8(8)	1062.3(1.7)	597.2(2.4)
1064	0.04281	270.8(8)	233.6(2)	597.5(1.2)	686.9(9)	1162.1(1.9)	627(3)
1060	0.04297	272.1(8)	234.3(2)	602.4(1.2)	693.1(9)	1180.0(1.9)	632(3)
1053	0.04326	274.6(8)	235.7(3)	611.3(1.2)	704.4(9)	1213.2(1.9)	641(3)
1047	0.04351	276.7(8)	237.0(3)	619.3(1.2)	714.5(9)	1243.8(2.0)	650(3)
1030	0.04422	283.2(8)	240.7(3)	643.9(1.3)	746.0(9)	1343.6(2.1)	675(3)
985	0.04624	304.0(8)	252.0(3)	728.3(1.4)	855.8(1.0)	1759(3)	765(3)
980	0.04648	306.7(8)	253.5(3)	739.9(1.4)	871.2(1.0)	1828(3)	778(3)
946	0.04815	327.8(9)	264.4(3)	836.4(1.6)	1001.6(1.1)	2581(4)	886(3)
930	0.04898	339.6(9)	270.3(3)	895.9(1.7)	1084.4(1.2)	3325(5)	957(3)
799.3	0.05699	549.7(1.1)	353.8(4)	3666(7)	13483(14)	-2333(4)	-1474(11)
700	0.06507	1983(3)	553.1(0.6)	-1394(3)	-1192.7(1.4)	-707.9(1.4)	-3787(18)
600	0.07592	-645.9(1.3)	4506(5)	-438.4(0.9)	-421.8(0.7)	-333.9(1.0)	-523(3)
500	0.09110	-200.1(0.8)	-411.1(0.4)	-201.3(0.5)	-196.8(0.6)	-166.3(1.0)	-204(2)
400	0.11388	-86.4(0.8)	-134.0(0.2)	-131.4(2.4)	-109.7(1.1)	-88.2(1.3)	-111(4)
300	0.15183	-38.4(1.1)	-57.5(0.3)	-47.2(0.4)	-43.7(1.1)	-35(2)	-43(2)

The calculation of the core polarizability is carried out in the random-phase approximation (RPA); we verified that our static values agree with RPA values from Ref. [10]. The separation of the polarizability into core and valence parts also requires the addition of the compensation term α_{vc} which accounts for the contribution from the excitation to the occupied valence shell that is forbidden by the Pauli exclusion principle. The calculation of the core polarizability does not exclude the excitation from the core to the valence shell and half of this contribution has to be subtracted. Thus, the polarizability contributions may be separated as

$$\alpha^{ns} = \alpha_{core} + \alpha_{vc} + \alpha_v^{ns}. \quad (1)$$

We calculate the valence part of the ac polarizability of the ground state ns for the alkali-metal atoms by computing the sum over states

$$\alpha_v^{ns} = \frac{1}{3} \sum_{n'} \left(\frac{\delta E_{n'p_{1/2}} \langle n'p_{1/2} || D || ns \rangle^2}{\delta E_{n'p_{1/2}}^2 - \omega^2} + \frac{\delta E_{n'p_{3/2}} \langle n'p_{3/2} || D || ns \rangle^2}{\delta E_{n'p_{3/2}}^2 - \omega^2} \right), \quad (2)$$

where $\langle np || D || ns \rangle$ is the reduced electric-dipole matrix element, $\delta E_{n'p_{1/2}} = E_{n'p_{1/2}} - E_{ns}$, and $\delta E_{n'p_{3/2}} = E_{n'p_{3/2}} - E_{ns}$. In this formula, ω is assumed to be at least several linewidths off resonance with the corresponding transition. We use the system of atomic units,

a.u., in which $e/\sqrt{4\pi\epsilon_0}$, m_e , and the reduced Planck constant $\hbar$ have the numerical value 1, in Eq.(2). Polarizability in a.u. has the dimensions of volume, and its numerical values presented here are thus measured in units of a_0^3 , where $a_0 \approx 0.052918$ nm is the Bohr radius. The atomic units for α can be converted to SI units via $\alpha/h [\text{Hz}/(\text{V/m})^2] = 2.48832 \times 10^{-8} \alpha [\text{a.u.}]$, where the conversion coefficient is $4\pi\epsilon_0 a_0^3/h$ and Planck constant h is factored out. The atomic unit of frequency ω is $E_h/\hbar \approx 4.1341 \times 10^{16}$ Hz, where E_h is the Hartree energy.

The sum over n' in Eq.(2) converges rapidly, and unless the frequency is resonant with the transitions other than the primary ones ($ns - np_{1/2}$ and $ns - np_{3/2}$), the first term is dominant. Therefore, only a few terms need to be calculated accurately. We use the experimental numbers compiled in Ref. [11] together with their uncertainties for the first term, for example for $n' = 3$ term for Na. High-precision theoretical all-order values are used for the next three terms (for example $n' = 4, 5, 6$ terms for Na). We refer the reader to Refs. [11, 12, 13] for the detailed description of the single-double (SD) all-order method and its extensions. The all-order values of the matrix elements are evaluated for their accuracy, and the extensions of the SD all-order method (SDpT, which partially includes triple excitations or SD_{sc} , which includes semi-empirical scaling of dominant terms) are used for certain transitions where those values are expected to be of better accuracy. The uncertainty of the resulting matrix elements is evaluated based on the relative impor-

TABLE II: The parameters A^{ns} (in a.u.) for the calculation of ground state frequency-dependent polarizabilities of alkali-metal atoms in α_0^3 for the wavelengths above 800 nm. The corresponding uncertainties in the values of A^{ns} are given in the last column. The corresponding reduced matrix elements and energy differences are given in columns labeled $\langle np||D||ns \rangle$ and δE_{np} .

Atom	n	$\langle np_{1/2} D ns \rangle$	$\langle np_{3/2} D ns \rangle$	$\delta E_{np_{1/2}}$	$\delta E_{np_{3/2}}$	A^{ns}	δA^{ns}
Li	2	3.317(4)	4.689(5)	0.067906	0.067907	2.04	0.69
Na	3	3.5246(23)	4.9838(34)	0.077258	0.077336	1.86	0.12
K	4	4.102(5)	5.800(8)	0.059165	0.059428	6.26	0.33
Rb	5	4.231(3)	5.977(4)	0.057314	0.058396	10.54	0.60
Cs	6	4.4890(65)	6.3238(73)	0.050932	0.053456	17.35	1.00
Fr	7	4.277(8)	5.898(15)	0.055758	0.063442	24.8	1.8

tance of certain classes of the all-order terms, extensive comparison of various atomic properties of alkali-metal atoms with experiment [9, 11, 13], and the spread of SD, SDpT, and SD_{sc} values. Some of these values were published previously in Refs. [11, 14]. The experimental values from [14] are used for $6s - 7p_{1/2}$ and $6s - 7p_{3/2}$ transitions. In summary, the set of values used for the calculation of the first four terms in the sum over states of Eq.(2) consists of the best known values for these transitions. The experimental energies from Refs. [15, 16] are used for the corresponding energy levels. We refer to the total contribution from the first four terms as the main term; the remaining contributions are referred to below as the tail contribution; i.e.

$$\alpha_v^{ns} = \alpha_{\text{main}}(n' = n, \dots, n+3) + \alpha_{\text{tail}}(n' > n+3). \quad (3)$$

We evaluated the tail contributions with a complete set of basis Dirac-Hartree-Fock (DHF) wave functions generated using the B-spline method [17]. We use 40 splines of order $k = 7$ for each angular momentum. The basis set orbitals are defined on a non-linear grid and are constrained to a large spherical cavity of a radius $R = 75 - 110$ a.u. The cavity radius is chosen to accommodate first four ns , $np_{1/2}$, and $np_{3/2}$ valence orbitals. For consistency, we use the same basis sets for the tail calculations as we used in the all-order calculation of the main term matrix elements. The tail contribution is calculated in the DHF approximation. We list our results for the frequency-dependent polarizabilities from the ultraviolet through the infrared spectrum in Table I. Particular attention is given to the values for the infrared wavelengths which have been or might be employed in the present generation of optical traps. Our values of the ac polarizabilities are accurate to better than 0.5% with the exception of the values corresponding to wavelengths in the ultraviolet spectrum. For these wavelengths, non-primary resonances become dominant and the uncertainty of the polarizability values is dominated by the uncertainty in the corresponding resonance matrix elements. For example, the first resonances in Rb ($5s - 5p$ transitions) occur at 795 nm and 780 nm, while next resonances ($5s - 6p$) transitions are near 420 nm. As the accuracy of the $5s - 6p$ matrix elements is lower than that of the $5s - 5p$ matrix elements the resulting accuracy




FIG. 1: The ratios $\alpha(\lambda)/m$ of frequency-dependent polarizabilities to the atomic weights for ${}^6\text{Li}$ and ${}^{23}\text{Na}$.

of the ac polarizabilities at 400 nm is also lower.

III. RESULTS

We have investigated the dependence on the frequency ω of all of the terms contributing to α^{ns} , where ns is the ground state. We find that the α_{core} , α_{vc} , α_{tail} , and α_v^{ns} terms with $n' = n+1, n+2, n+3$ depend weakly on the value of ω for wavelengths above 800 nm. In fact, it is possible to accurately parameterize the values of the ac polarizabilities for those wavelengths by the formula

$$\alpha_v^{ns}(\omega) = \frac{1}{3} \frac{\delta E_{np_{1/2}} \langle np_{1/2}||D||ns \rangle^2}{\delta E_{np_{1/2}}^2 - \omega^2} + \frac{1}{3} \frac{\delta E_{np_{3/2}} \langle np_{3/2}||D||ns \rangle^2}{\delta E_{np_{3/2}}^2 - \omega^2} + A^{ns}, \quad (4)$$

where A^{ns} is independent on the value of ω . All values in Eq.(4) are in atomic units. The values of the constants A^{ns} and their uncertainties are given in Table II. The values of A^{ns} are calculated as the aver-

TABLE III: Selected wavelengths λ_{air} in nm, at which two species of optically-trapped alkali atoms (^6Li , ^7Li , ^{23}Na , ^{40}K , ^{41}K , ^{87}Rb , and ^{133}Cs) would have the same frequencies of oscillation in a common optical trap. The corresponding ground state frequency-dependent polarizabilities of the alkali-metal atoms (in a.u.) are also given.

λ_{air}	1	α_1	2	α_2
557.56(13)	^6Li	-360(1)	^{23}Na	-1375(1)
768.648(3)	^6Li	681.3(1.3)	^{40}K	4525(112)
799.46(12)	^6Li	549(1)	^{40}K	3649(7)
768.645(3)	^6Li	681.3(1.3)	^{41}K	4638(112)
798.44(12)	^6Li	553(1)	^{41}K	3763(7)
786.07(1)	^6Li	597.9(1.2)	^{87}Rb	8639(20)
804.57(4)	^6Li	533(1)	^{87}Rb	7704(8)
862.097(28)	^6Li	412.5(1.0)	^{133}Cs	9113(22)
904.76(4)	^6Li	361.7(9)	^{133}Cs	7991(12)
549.72(18)	^7Li	-329(1)	^{23}Na	-1078(1)
768.665(3)	^7Li	681(1)	^{40}K	3880(112)
806.9(2)	^7Li	526(1)	^{40}K	2998(6)
768.662(3)	^7Li	681(1)	^{41}K	3978(112)
805.55(15)	^7Li	530(1)	^{41}K	3096(6)
786.56(1)	^7Li	595.9(1.2)	^{87}Rb	7383(19)
807.16(4)	^7Li	526(1)	^{87}Rb	6510(6)
863.436(32)	^7Li	410.5(1.0)	^{133}Cs	7777(20)
907.25(6)	^7Li	359.3(9)	^{133}Cs	6805(10)
789.240(7)	^{23}Na	364.9(4)	^{87}Rb	1380(16)
946.664(5)	^{23}Na	264.1(3)	^{87}Rb	999(1)
875.53(3)	^{23}Na	295.5(3)	^{133}Cs	1709(11)
1021.5(4)	^{23}Na	242.6(3)	^{133}Cs	1403(2)
768.848(3)	^{40}K	-3893(116)	^{87}Rb	-8470(10)
784.32(1)	^{40}K	6806(13)	^{87}Rb	14800(27)
807.31(6)	^{40}K	2968(6)	^{87}Rb	6454(6)
868.90(4)	^{40}K	1298(3)	^{133}Cs	4317(14)
938.6(2)	^{40}K	862(2)	^{133}Cs	2868(4)
768.851(3)	^{41}K	-4013(116)	^{87}Rb	-8472(10)
784.42(1)	^{41}K	6767(12)	^{87}Rb	14358(26)
808.07(7)	^{41}K	2916(6)	^{87}Rb	6187(6)
869.14(4)	^{41}K	1296(2)	^{133}Cs	4203(14)
940.9(2)	^{41}K	854(2)	^{133}Cs	2771(4)
790.303(6)	^{87}Rb	-1299(17)	^{133}Cs	-1986(3)
873.39(4)	^{87}Rb	1624(2)	^{133}Cs	2483(12)
1158(3)	^{87}Rb	580.5(8)	^{133}Cs	888(2)

age values of the totals of the α_{core} , α_{vc} , α_{tail} , and $\alpha_{\text{v}}^{n'}$ terms with $n' = n + 1, n + 2, n + 3$ calculated in the range from 800 nm to 1600 nm. The values of the $\langle np || D || ns \rangle$ reduced electric-dipole matrix elements and the corresponding energy differences δE are also given. The electric-dipole matrix elements for Li are experimental values from Ref. [18], all other matrix elements are experimental values compiled in [11]. The energy differences are taken from Refs. [15, 16]. The values of the ac polarizabilities obtained using Eq.(4) differ from our results in Table I in the relevant wavelength range by less




FIG. 2: The ratios $\alpha(\lambda)/m$ of frequency-dependent polarizabilities to the atomic weights for ^{40}K and ^{87}Rb .

than 0.1% which is significantly below the accuracy of the values themselves. We note that the formula (4) is not applicable for the lower wavelengths owing to the importance of the other $n'p - ns$ resonances. For example, the 420 nm wavelength corresponds to $6p - 5s$ transition in Rb; therefore, the ground state polarizability at this frequency will be dominated by the $n' = 6$ term.

IV. THE FREQUENCY-MATCHING CRITERIA FOR MIXED SPECIES

We have also conducted a search of the wavelength values at which two species of optically-trapped alkali atoms (^6Li , ^7Li , ^{23}Na , ^{40}K , ^{41}K , ^{87}Rb , ^{133}Cs , and ^{210}Fr) would have the same frequencies of oscillation in a common optical trap, i.e. λ such that

$$s \equiv \sqrt{\frac{\alpha_1(\lambda)}{m_1} \frac{m_2}{\alpha_2(\lambda)}} \approx 1, \quad (5)$$

where m_i is the atomic weight [19]. The wavelengths of those selected matches and their uncertainties are listed in Table III. We have omitted all matches at wavelengths below 500 nm. While we had found numerous matches in the range from 300 nm to 500 nm they are difficult to place accurately since the accuracy of our calculation is lower in this region. The width of the matches in the 300 – 500 nm region is generally very narrow and the corresponding values of the polarizabilities are small, making them of limited experimental use. We define the width of the match as the wavelength range where the parameter s given by Eq. (5) is between 0.9 and 1.1. The width of the match is very narrow if the corresponding wavelength is in the immediate vicinity of the resonance. We have also omitted most matches at higher wavelengths from Table III that are close to the resonance.

Note that we listed the values of the wavelengths in the air. While the difference between air and vacuum wavelengths is small in the considered region, 0.15–0.4 nm, it is sufficient to cause changes in the frequency-dependent polarizabilities (at some wavelengths) which exceed the uncertainties of the calculations. We used the following formula to compute the wavelength in air from the vacuum wavelength[20]

$$\lambda_{\text{air}} = \lambda_v / (1 + 10^{-8} (8060.51 + \frac{2480990}{132.274 - \sigma^2 \times 10^6} + \frac{17455.7}{39.32957 - \sigma^2 \times 10^6})). \quad (6)$$

The wavenumber σ is here to be expressed in reciprocal nm. We also list the values of the ac polarizabilities and their uncertainties in Table III.

The uncertainty in the wavelength of the match results from the uncertainties in the values of the frequency-dependent polarizabilities $\alpha_1(\lambda)$ and $\alpha_2(\lambda)$ given in Table III. The uncertainties in the values of $\alpha(\lambda)$ are found to be independent of λ within the range of the match to a very good precision but are generally different for α_1 and α_2 . We calculate the resulting uncertainties $\delta\lambda_1$ and $\delta\lambda_2$ due to uncertainties in the values of α_1 and α_2 separately and add the results in quadrature to obtain the final uncertainty $\delta\lambda$. To evaluate $\delta\lambda_1$, we calculate the $\alpha_1^\pm(\omega)/m_1 = [\alpha_1(\lambda) \pm \delta\alpha_1]/m_1$ and then determine the wavelengths $\lambda^\pm$ at which $\alpha_1^\pm(\lambda)/m_1 = \alpha_2(\lambda)/m_2$. The maximum of the differences between the initial match wavelength and $\lambda^\pm$ is taken to be the uncertainty $\delta\lambda_1$. The evaluation of $\delta\lambda_2$ is done in the same way.

We illustrate several of the matches given in Table III in Fig. 1 and Fig. 2. The ratios $\alpha(\lambda)/m$ of frequency-dependent polarizabilities to the atomic weights for Li and Na in the wavelength range 500–800 nm are plotted in Fig. 1. The corresponding matches are shown by the arrows and the corresponding wavelengths are given on

the plot. The values $\frac{\alpha}{m}$ for ^6Li and ^{23}Na match at three wavelengths in the wavelength range plotted on Fig. 1. Two of the matches, at 589.397 nm and 670.79 nm, are very close to the resonances and, therefore, are not listed in Table III as discussed before. The match areas for those cases are extremely narrow and the polarizability of the corresponding alkali is very sensitive to very small changes in the wavelength. Fig. 2 shows similar plot to illustrate the wavelengths at which ^{40}K and ^{87}Rb have the same frequencies of oscillation in a common optical trap in the range 740–820 nm.

V. CONCLUSIONS

In summary, we conducted a systematic study of the ground state frequency-dependent polarizabilities of the alkali atoms from ultraviolet through infrared spectral regions. The values of the ac polarizabilities and their uncertainties are calculated for a number of wavelengths, including the wavelengths of common infrared lasers. A combination of high-precision measurements of the ratios of frequency-dependent polarizabilities of different species could provide excellent tests of the current experimental and theoretical values of the electric-dipole matrix elements in alkali-metal atoms. We provide formulas and the necessary parameters for the accurate calculation of the ac polarizabilities for all alkali-metal atoms at wavelengths above 800 nm. Finally, we list selected wavelengths at which two species of alkali atoms would have the same oscillation frequencies in a common optical trap.

This work was performed under the sponsorship of the U.S. Department of Commerce, National Institute of Standards and Technology.

-
- [1] N. Lundblad, D. C. Aveline, R. J. Thompson, J. M. Koehl, J. Ramirez-Serrano, W. M. Klipstein, D. G. Enzer, N. Yu, and L. Maleki, *J. Opt. Soc. Am. B* **21**, 3 (2004).
 - [2] F. Schreck, L. Khaykovich, K. L. Corwin, G. Ferrari, T. Bourdel, J. Cubizolles, and C. Salomon, *Phys. Rev. Lett.* **87**, 080403 (2001).
 - [3] Z. Hadzibabic, C. A. Stan, K. Dieckmann, S. Gupta, M. W. Zwierlein, A. Gorlitz, and W. Ketterle, *Phys. Rev. Lett.* **88**, 160401 (2002).
 - [4] G. Modugno, G. Roati, F. Riboli, F. Ferlaino, R. J. Brecha, and M. Inguscio, *Science* **297**, 2240 (2002).
 - [5] C. A. Stan, M. W. Zwierlein, C. H. Schunck, S. M. F. Raupach, and W. Ketterle (2004), *cond-mat/0406129*.
 - [6] S. Inouye, J. Goldwin, M. L. Olsen, C. Ticknor, J. L. Bohn, and D. S. Jin (2004), *cond-mat/0406208*.
 - [7] M. A. Morales, N. Nygaard, J. E. Williams, and C. W. Clark, *New J. Phys.* **7**, 87 (2005).
 - [8] M. S. Safronova, C. J. Williams, and C. W. Clark, *Phys. Rev. A* **67**, 040303(R) (2003).
 - [9] M. S. Safronova, C. J. Williams, and C. W. Clark, *Phys. Rev. A* **69**, 022509 (2004).
 - [10] W. R. Johnson, D. Kolb, and K.-N. Huang, *At. Data Nucl. Data Tables* **28**, 333 (1983).
 - [11] M. S. Safronova, W. R. Johnson, and A. Derevianko, *Phys. Rev. A* **60**, 4476 (1999).
 - [12] S. A. Blundell, W. R. Johnson, and J. Sapirstein, *Phys. Rev. A* **43**, 3407 (1991).
 - [13] M. S. Safronova, A. Derevianko, and W. R. Johnson, *Phys. Rev. A* **58**, 1016 (1998).
 - [14] A. A. Vasilyev, I. M. Savukov, M. S. Safronova, and H. G. Berry, *Phys. Rev. A* **66**, 020101(R) (2002).
 - [15] C. E. Moore, *Atomic Energy Levels*, vol. 35 of *Natl. Bur. Stand. Ref. Data Ser.* (U.S. GPO, Washington, D.C., 1971).
 - [16] J. R. Fuhr, W. C. Martin, A. Musgrove, J. Sugar, and W. L. Wiese, *NIST Atomic Spectroscopic Database*,

- NIST Physical Reference Data, National Institute of Standards and Technology. Available online, URL = http://physics.nist.gov/cgi-bin/AtData/main_asd.
- [17] W. R. Johnson, S. A. Blundell, and J. Sapirstein, Phys. Rev. A **37**, 307 (1988).
- [18] W. R. Johnson, Z. W. Lui, and J. Sapirstein, At. Data and Nucl. Data Tables **64**, 279 (1996).
- [19] J. S. Coursey, D. J. Schwab, and R. A. Dragoset, *NIST Nuclear Physics Database*, NIST Physical Reference Data, National Institute of Standards and Technology. Available online, URL = <http://physics.nist.gov/PhysRefData/Compositions/index>.
- [20] E. Peck and K. Reeder, J. Opt. Soc. Am. **63**, 958 (1972).